

Brissi in cornered skin and should have
liked to have had the advantage of seeing
more of them. I met them for a few
minutes on the platform at Taunton
and hoped to have travelled with them
as far as Bristol but the compartments
into which they got was full and I was
obliged to take another and had no time
even to take leave of them - They then
left when I returned to town

Your very sincerely
George Bentham

25. WILTON PLACE.
London
Oct 1/78

My dear Gray

I have to thank you for your
lecture on forest distribution which I am
much pleased with - It is a very lucid
exposition of the theory you have so well
worked out of the origin and history of
North American forest vegetation. We
want much a similar clue to the
distribution of tropical trees but unfortunately
have very much fewer data to go upon
than is the case with the northern ancient
and modern trees

Much obliged for your flattering
article on my Australian flora - it has
not been noticed as far as I know in any

European journals - I forgot to insert in
the Preface as I intended that the total
number of species described is 8168 in
1600 genera. Mueller thinks he has
about 500 species already to add. I wish
I could have fulfilled my original plan
of a supplement and review of the
geographical distribution etc but I have
just passed my 78th birthday and am
obliged to devote what little time I may
still have before me to genera Plantarum.
I have not quite done yet with Euphorbiaceae
which have proved a tougher job than I
thought - I made 6 tribes Euphorbiae,
Pleurolobiae, Baccin., Phyllanthae, Galerinae
and Cootoneae - Phyllanthae with the

outer stamens when numerous opposite
the raphe and cells of the ovary 2-ovulate
Cootoneae with the outer stamens when
numerous alternate with the raphe and
cells of the ovary 1-ovulate - the Galerinae
a few genera only with the stamens of
Phyllanthae and the ovary of Cootoneae
obtusiflora exorosa etc included in
Phyllanthae

Can you let me have female
flowers (not yet grown into fruits) of
Sinuomordia - I do not understand how
this genus can have been put into
Baccin. but our specimens are either
males in flower or females in fruit.

I had great pleasure in meeting
Dr. W. James the other day at Dr.